

Decaying Λ cosmology, varying G and holography

S. Carneiro¹ and J. A. S. Lima^{2,3}

¹*Instituto de Física, Universidade Federal da Bahia, 40210-340, Salvador, BA, Brazil*

²*Departamento de Astronomia - IAG, Universidade de São Paulo, 05508-900, São Paulo, SP, Brazil*

³*Departamento de Física, Universidade Federal do Rio Grande do Norte, 59072-970, Natal, RN, Brazil*

We discuss a class of uniform and isotropic, spatially flat, decaying Λ cosmologies, in the realm of a model where the gravitation “constant” G is a function of the cosmological time. Besides the usual de Sitter solution, the models at late times are characterized by a constant ratio between the matter and total energy densities. One of them presents a coasting expansion where the matter density parameter is $\Omega_m = 1/3$, and the age of the universe satisfies $Ht = 1$. From considerations in line with the holographic conjecture, it is argued that, among the non-decelerating solutions, the coasting expansion is the only acceptable from a thermodynamic viewpoint, and that the time varying cosmological term must be proportional to H^2 , a result earlier obtained using different arguments.

I. INTRODUCTION

One of the most intriguing problems in modern cosmology is the present observed value of the cosmological constant, which amounts to nearly 120 orders of magnitude smaller than the value predicted by quantum field theories [1]. A possible explanation for this huge discrepancy is based on the idea that the vacuum energy density is not constant, but decays as the universe expands [2]-[14].

When the vacuum energy density is computed with the help of quantum field theories in flat spacetime, one obtains a divergent result, which can be regulated by means of an ultraviolet cutoff of the order of the Planck mass. In the standard recipe, such a contribution to the vacuum energy must be exactly canceled by a bare cosmological constant in the Einstein equations, because, in the flat spacetime, the right-hand side of those equations is identically zero [13]. Therefore, in a curved background, a renormalized value for Λ should be obtained by calculating the vacuum energy density and then subtracting the Minkowskian result. Since spacetime was strongly curved at Planck times, one may expect an initial huge value of the order of l_p^{-2} , where $l_p = \sqrt{G}$ is the Planck length. But, as the universe expands, the observed cosmological constant should decay, thereby leading to the small value observed nowadays. In other words, the present value of Λ (the vacuum energy density) is small because the Universe is too old [4].

This line of reasoning leads one to infer that other physical “constants” should also vary at a cosmological scale. Indeed, the quantum vacuum fluctuations behind the net vacuum energy density contribute also to the observed, renormalized, values of couplings and masses of elementary particles. Therefore, if the vacuum density decays with the universe expansion, one should expect cosmological time variations of charges and masses. The first would lead to the variation of the fine structure constant [15], an effect that have possibly been observed [16]. As to the variation of masses, it is equivalent to a variation of the gravitational constant G , at least in what

concerns gravitational effects. The variation of G should also be expected by considering it a renormalized coupling constant of a (still unknown) quantum theory of gravity.

Since the seminal papers by Dirac [17], a possible variation of G has been investigated with no success by several teams, through geophysical and astronomical observations, at the scale of solar system and with binary systems [18]. However, it should be stressed that we are talking here about time variations at a cosmological scale, and cosmological observations still cannot put strong limits on such a variation, specially at the late times of the evolution. Naturally, if the G variation originates from the decaying of vacuum density at a cosmological scale, one would expect a constant value of G at the scales of solar system and galaxies, since at these scales the matter density and the curvature (and presumably the vacuum density) are nearly constant.

In this paper we discuss a new decaying vacuum scenario which is supposed to induce a time varying gravitational constant. As we shall see, the Friedmann equations for the flat case plus the Eddington-Weinberg relation [19] imply that beyond the usual de Sitter solution there are other solutions characterized by a constant ratio between the matter density and the total energy density. One of them presents a coasting expansion where the universe age is fixed by $Ht = 1$, and the matter density parameter is $\Omega_m = 1/3$.

The work is structured as follows. Next section we review an earlier approach to decaying vacuum energy models with varying G [7, 10]. In the subsequent sections we make some considerations in line with the so called holographic principle [11, 19, 20] to justify the Eddington-Weinberg relation, and to argue that, among all the possible non-decelerating solutions, that with coasting expansion is the only plausible from a thermodynamic viewpoint.

It should be noticed that models with variation of both Λ and G have already been considered in the literature. For instance, Bertolami [3] gave a detailed analysis of varying Λ cosmological solutions in the realm of a Brans-Dicke theory. However, the variation of G we are inter-

ested here has its origin in the variation of the vacuum energy density. Therefore, although considering that a Brans-Dicke scalar field might be used as an effective description of that variation, in what follows we shall make no use of such a field.

II. DECAYING Λ SOLUTIONS

In the flat case, the Friedman equations with a time dependent G can be written as

$$\rho = \frac{3H^2}{8\pi G}, \quad (1)$$

$$\dot{\rho} + 3H(\rho + p) = 0, \quad (2)$$

where $H = \dot{a}/a$ is the Hubble parameter, and a dot means derivation respect to the cosmological time t . The second equation does not depend on the varying or constant character of G , being just an expression of energy conservation and the equivalence principle [7, 12]. As far as relation (1) is concerned, it is valid today, and we will assume that it is also valid for any time in course of the evolution.

The variation law for G follows from the Eddington-Weinberg empirical relation [19]

$$G = \frac{H}{8\pi\lambda} \approx \frac{H}{m^3}, \quad (3)$$

where m has the order of the pion mass, and the constant λ was introduced for convenience. Once again, this relation is valid today, and we will assume that it is valid since early times. Substituting (3) into (1), we have

$$\rho = 3\lambda H. \quad (4)$$

Let us now consider a twofold energy content, formed by dust matter with energy density ρ_m plus a vacuum term with equation of state $p_\Lambda = -\rho_\Lambda$. The total energy density and pressure read

$$\rho = \rho_m + \rho_\Lambda, \quad (5)$$

$$p = -\rho_\Lambda. \quad (6)$$

Inserting (4)-(6) into the conservation equation (2), we obtain

$$\lambda\dot{H} + \rho_m H = 0. \quad (7)$$

To proceed further, let us now assume that the quantities H and ρ_m , at late times, fall monotonically as power laws of the scale factor a :

$$H = \beta/a^k, \quad (8)$$

$$\rho_m = \gamma/a^n, \quad (9)$$

where n and k are two positive parameters. Replacing the above expressions into (7) leads to

$$\gamma a^{-n} - \lambda\beta k a^{-k} = 0. \quad (10)$$

The above equation is valid for any (large) value of a only if

$$k = n, \quad (11)$$

$$\gamma = \lambda\beta n. \quad (12)$$

With these results, and by using (4), (8) and (9), it is easy to show that

$$\Omega_m \equiv \rho_m/\rho = n/3. \quad (13)$$

Since $0 \leq \Omega_m \leq 1$, it follows that $0 \leq n \leq 3$. In the case $n = 0$, we have

$$H = \beta, \quad (14)$$

$$\rho_m = 0, \quad (15)$$

$$G = \frac{\beta}{8\pi\lambda}. \quad (16)$$

This solution corresponds to a de Sitter universe, with Λ and G constant.

For the other values of n , we obtain from (8) the solution [21]

$$a = (n\beta t)^{1/n}, \quad (17)$$

$$Ht = 1/n, \quad (18)$$

$$q = n - 1, \quad (19)$$

where $q = -a\ddot{a}/\dot{a}^2$ is the deceleration parameter. In the case $n = 1$ we see that $q = 0$ and

$$a = \beta t, \quad (20)$$

$$Ht = 1, \quad (21)$$

$$\Omega_m = 1/3. \quad (22)$$

As discussed in [22], among the solutions of the form $a \propto t^n$, the best fitting of supernova and radio sources observations is obtained in the coasting case, $a \propto t$. On the other hand, the relation $Ht = 1$ gives the best estimation for the universe age [23], and the relative matter density given by (22) matches surprisingly well the observed value [24].

It is worth notice that in a context with varying Λ but constant G the coasting solution with $Ht = 1$ corresponds to a relative matter density equals to $2/3$ [10], in clear disagreement with observation. To obtain $\Omega_m = 1/3$, we would have to have $Ht = 2$, completely outside the observed limits.

Let us determine the variation rate of G , as well as the rate of matter production in this model. The evolution law (3) leads to the relative variation rate

$$\dot{G}/G = -(1 + q)H. \quad (23)$$

For $n = 1$, we have $q = 0$, and so

$$\dot{G}/G = -H. \quad (24)$$

The rate of matter production (coming from the decaying vacuum energy) reads

$$\frac{1}{\rho_m a^3} \frac{d}{dt} (\rho_m a^3) = (3 - n)H. \quad (25)$$

Let us now discuss the variation of the vacuum energy density in the present context. By using (3)-(5) and (13), one may check that the vacuum energy density and the corresponding cosmological constant are given by

$$\rho_\Lambda = (3 - n)\lambda H \approx \frac{(3 - n)}{8\pi} m^3 H, \quad (26)$$

$$\Lambda = 8\pi G \rho_\Lambda = (3 - n)H^2, \quad (27)$$

leading to present values in agreement with observation.

The first above equation is in accordance with a recent derivation by Schützhold [13], based on quantum field estimations in an expanding background. The second equation agrees with an ansatz proposed by Chen and Wu on the basis of dimensional arguments [5], and further modified by Carvalho et al. [6] in order to obtain the H^2 dependence. It also matches a result based on a renormalization approach [14], in the spirit discussed in the Introduction. Note that, in the realm of a constant G cosmology, those two equations would be incompatible to each other [25].

III. HOLOGRAPHY AND G VARIATION

Since the initial proposals on the existence of entropy bounds in self-gravitating systems, rigorous verifications of this idea have been obtained in some cases, leading to the general belief that there is an entropy bound associated to any horizon, of the order of the horizon area (in Planck units) [20]. For our purposes, this holographic conjecture could be understood as follows.

Consider a set of degrees of freedom inside a horizon of characteristic radius R . The limitation of its configuration space leads to the quantization of its energy-momentum space, with a quantum of energy given by $m_0 \approx 1/R$ [26]. Then, the maximum number of degrees of freedom inside the horizon is given by $N \approx M/m_0 \approx MR$, where M is the mass inside the horizon. So, for any system obeying the relation $M \approx R/G$, we have $N \approx R^2/G$, as just established by the holographic conjecture.

The reader knows two important examples of systems satisfying the above conditions. The first is a black-hole, for which we have (in the non-rotating case) $M = R/(2G)$, where R is the gravitational radius. The second is the spatially flat Friedmann universe. Indeed, in this case the total energy density is given by $\rho = 3/(8\pi G R^2)$, where R is the Hubble radius. On the other hand, the volume inside the Hubble horizon is $4\pi R^3/3$, which leads to a total mass inside the Hubble sphere given, again, by $M = R/(2G)$. Therefore, the holographic conjecture applies as well to our universe: the number of observable

degrees of freedom is bounded by the area of the Hubble horizon.

But how many degrees of freedom we have in the observable universe? We know that the number of barions is of the order of 10^{80} , and that the Universe contains about 10^8 photons per barion. It is also reasonable to believe that the contribution of the non-barionic dark matter may be of the same order of magnitude. But the major contribution to the entropy of matter seems to come from massive black-holes present in galactic nuclei, which represents an entropy of the order of 10^{101} [27].

As far as the vacuum entropy is concerned, it is not trivial to define the number of virtual particles in curved backgrounds. Let us remind, however, that our universe has a quasi-flat spacetime. Therefore, the following estimation can be considered a good approximation [11]. It is clear that vacuum configurations of classical fields (as the vacuum expectation value of the Higgs field or the QCD condensates) do not contribute to the vacuum entropy. In what concerns the zero-point fluctuations, they have, properly speaking, an infinite entropy density, because (if we do not impose any energy cutoff) the number of modes is infinite. But if we regulate their energy, by introducing an ultraviolet cutoff m , we also regulate their entropy. A simple estimation of the resulting entropy bound can be derived as follows.

Limiting the energy-momentum space associated to the zero-point fluctuations leads to the quantization of their configuration space, with a minimum size given by $l \approx m^{-1}$. This results in a superior bound to the number of available degrees of freedom in a given volume, say, the volume inside the Hubble horizon. The maximum number N_{max} of observable degrees of freedom will be of the order of V/l^3 , where V is the Hubble volume. That is,

$$N_{max} \approx \left(\frac{R}{l}\right)^3 \approx \left(\frac{m}{H}\right)^3. \quad (28)$$

Note that this superior bound dominates over the matter entropy. Indeed, taking for H the value observed nowadays, $H \approx 70 \text{ km.s}^{-1} \text{ Mpc}^{-1}$, we obtain from (28) (with m equals to the pion mass, see below) $N_{max} \approx 10^{122}$, a value that predominates over the matter entropy referred above, of order 10^{101} . Since N_{max} increases with R^3 , this dominance will remain valid in the future, and we can consider the entropy bound of the observable universe as given by (28).

On the other hand, we have seen that this bound should be equal to

$$N \approx \left(\frac{R}{l_p}\right)^2 = (H l_p)^{-2}. \quad (29)$$

Therefore, we can identify (29) with (28), which leads to equation (3), that is, to the Eddington-Weinberg relation.

But why has m the order of the pion mass? The common belief is that a natural cutoff for the vacuum fluctuations should be given by the Planck mass $m_p = l_p^{-1}$,

because at the Planck scale the classical picture of space-time breaks down. However, it is not difficult to see that, equating (29) to (28) with $m = m_p$, one obtains a Hubble radius of the order of l_p , which is not consistent with our universe.

One can also argue that the fluctuations of other fields than quarks and gluons contribute to the vacuum entropy as well. It is then intriguing that just the pion mass enters in the Eddington-Weinberg relation. Nevertheless, let us remark that in a curved spacetime the different sectors of the Standard Model of particles interactions are coupled by gravity. Then one may expect that (at late times) all the interacting vacuum fields tend to a state of thermodynamic equilibrium, at a temperature equal or below the temperature of the last vacuum phase transition. But the last of such transitions was the chiral transition of QCD [28], at a temperature given by $\Lambda_{QCD} \approx 150$ MeV, of the order of the pion mass.

IV. HOLOGRAPHY AND DECAYING Λ SOLUTIONS

As we have seen, the energy inside the Hubble horizon of the spatially flat Friedmann universe is given by

$$E = \frac{R}{2G} = 4\pi\lambda R^2 \quad (30)$$

(where we have used (3)). Therefore, when the Hubble radius has an increment dR the energy content increases by

$$dE = 8\pi\lambda R dR = dR/G. \quad (31)$$

On the other hand, in the case of a universe composed by dust and a vacuum term with pressure $p = -\Lambda/(8\pi G)$, as in (5)-(6), the general thermodynamic expression for the energy variation is given by

$$dE = TdS - p dV = Td\left(\frac{A}{\alpha G}\right) + \frac{\Lambda}{8\pi G} dV, \quad (32)$$

where the holographic prescription $S = A/(\alpha G)$ has been used (the positive dimensionless constant α is equal to 4 in the black-hole case). Equating (32) to (31), and using (3), $A = 4\pi R^2$, and $V = 4\pi R^3/3$, we obtain

$$T = \frac{\alpha H}{12\pi} \left(1 - \frac{\Lambda}{2H^2}\right). \quad (33)$$

This last result has important implications in relation to the cosmological solutions we have found in section II. First of all, we see that (in the case of a non-negative cosmological constant) $T \leq \alpha H/(12\pi)$. As far as the universe expands, the temperature decreases, and we have two different possibilities. The first one is that the universe tends asymptotically to a de Sitter universe, with constant Hubble radius and constant temperature (this

corresponds to the late time solution with $n = 0$ of section II). But in a de Sitter universe the Hubble constant is given by $H = \sqrt{\Lambda/3}$. Therefore, from (33) we have $T = -\alpha H/(24\pi)$. Since α is a positive constant, it's clear that such a solution is not acceptable from a thermodynamic point of view.

The second possibility is that the Hubble sphere expands forever, with $R \rightarrow \infty$ (our late time solutions with positive n). Leading (27) into (33), we obtain

$$T = (n-1) \frac{\alpha H}{24\pi}. \quad (34)$$

In this case, the temperature tends asymptotically to zero as far as the universe tends to a flat spacetime. But the condition $T \geq 0$ imposes $n \geq 1$. That is, the accelerating solutions (including the case $n = 0$, already discussed) are forbidden from this thermodynamic viewpoint. On the other hand, the late time coasting solution ($n = 1$) corresponds to $T/H = 0$. In other words, in the coasting expansion the temperature tends to zero faster than in the decelerating cases.

If the above analysis is repeated in the context of a model with varying Λ but constant G , the same conclusion is obtained: among the non-decelerating solutions, the coasting expansion is the only acceptable from a holographic perspective. However, in this case we would not be in accordance with the observed relative matter density, as already discussed. Note also that the holographic expression (33) suggests naturally the H^2 dependence for Λ discussed by several authors in different contexts [6, 14].

V. CONCLUSIONS

The asymptotic scenario proposed here has several interesting features, from both theoretical and observational viewpoints. A coasting expansion seems to provide the best fit of the recent observations of supernovas and radio sources [22]. Furthermore, it leads to a universe age given by $Ht = 1$, once again in good accordance with the observational limits [23]. Finally, a relative matter density equals to 1/3 is in remarkable agreement with observations [24], specially if we remember that it is an exact result in our context, and, perhaps more important, it does not depend on any fine tuning of model parameters or initial conditions.

From a theoretical perspective, the variation of the vacuum energy density with the cosmic expansion is naturally expected on the basis of quantum field theories in curved backgrounds [13, 14], as well as the association of a temperature to the vacuum of a curved spacetime. As discussed in the Introduction, the same can be said about the variation of the gravitational constant (and of the fine structure constant). The empirical law initially used for the variation of G (the Eddington-Weinberg relation, expression of Dirac's large numbers coincidence [7, 19]) can be justified on the basis of the holographic principle (see

also [11, 19]). Moreover, the late time coasting expansion is, among the non-decelerating solutions, the only compatible with a holographic, thermodynamic analysis.

Probably, it is not necessary to say that our analysis cannot be considered complete. First of all, the holographic idea we have adopted to justify the Eddington-Weinberg relation is still just a conjecture, despite the several theoretical developments this subject has received in the last years [20]. On the other hand, the Friedmann equation (1) applied to this varying G context is something to be theoretically validated, for example in the realm of scalar-tensor theories [3, 29]. It is possible to verify that our solutions, together with equation (1), are solutions of Brans-Dicke theories, and an exhaustive study about this will be shown elsewhere.

Finally, let us remind that our discussion is restricted

to the late eras of the expanding Universe. The study of a complete scenario, including the hot phase at early times, depends on the establishment of exact variation laws for Λ and G , which will depend on the development of a field theoretical treatment of the problem in a curved, expanding background. Such a study should also explain why the space is nearly flat, perhaps by means of a fundamental justification of inflation which may ultimately be related to a definite decaying law for Λ at very short times.

The authors are thankful to R. Abramo, A. I. Arbab, O. Bertolami, M. V. John, T. Mongan, T. Padmanabhan, A. Saa, J. Solà, J.-P. Uzan, and R. G. Vishwakarma for useful discussions and bibliographic suggestions. This work was partially supported by CNPq.

-
- [1] V. Sahni and A. Starobinsky, *Int. J. Mod. Phys. D* **9**, 373 (2000); P. J. E. Peebles and B. Ratra, *Rev. Mod. Phys.* **75**, 559 (2003).
 - [2] M. Ozer and O. Taha, *Phys. Lett. A* **171**, 363 (1986); *Nucl. Phys. B* **287**, 776 (1987).
 - [3] O. Bertolami, *Nuovo Cimento* **93**, 36 (1986).
 - [4] P. J. E. Peebles and B. Ratra, *Astrophys. J.* **325**, L17 (1988); B. Ratra and P. J. E. Peebles, *Phys. Rev. D* **37**, 3406 (1988).
 - [5] W. Chen and Y.S. Wu, *Phys. Rev. D* **41**, 695 (1990).
 - [6] J. C. Carvalho, J. A. S. Lima and I. Waga, *Phys. Rev. D* **46**, 2404 (1992).
 - [7] J. A. S. Lima and J. C. Carvalho, *Gen. Rel. Grav.* **26**, 909 (1994).
 - [8] J. A. S. Lima and J. M. F. Maia, *Phys. Rev. D* **49**, 5597(1994); J. A. S. Lima and M. Trodden, *Phys. Rev. D* **53**, 4280 (1996); J. A. S. Lima, J. M. F. Maia and N. Pires, *IAU Symposium* **198**, 111 (2000); J. V. Cunha, J. A. S. Lima and N. Pires, *Astron. and Astrophys.* **390**, 809 (2002).
 - [9] M. V. John and K. B. Joseph, *Phys. Rev. D* **61**, 087304 (2000); M. Novello, J. Barcelos-Neto and J. M. Salim, *Class. Quant. Grav.* **18**, 1261 (2001); **19**, 3107 (2002); R. G. Vishwakarma, *Class. Quant. Grav.* **18**, 1159 (2001); **19**, 4747 (2002); J. S. Alcaniz and J. M. F. Maia, *Phys. Rev. D* **67**, 043502 (2003); A. I. Arbab, *Class. Quant. Grav.* **20**, 93 (2003).
 - [10] S. Carneiro, gr-qc/0307114, Tenth Marcel Grossmann Meeting, Rio de Janeiro, 2003.
 - [11] S. Carneiro, *Int. J. Mod. Phys. D* **12**, 1669 (2003); gr-qc/0206064.
 - [12] R. Aldrovandi, J. P. Beltrán and J. G. Pereira, gr-qc/0312017.
 - [13] R. Schützhold, *Phys. Rev. Lett.* **89**, 081302 (2002); *Int. J. Mod. Phys. A* **17**, 4359 (2002).
 - [14] I. L. Shapiro and J. Solà, astro-ph/0401015.
 - [15] A. F. Rañada, *Europhys. Lett.* **61**, 174 (2003).
 - [16] J. K. Webb *et al.*, *Phys. Rev. Lett.* **87**, 091301 (2001).
 - [17] P. A. M. Dirac, *Nature* **139**, 323 (1937); *Proc. Roy. Soc. Lond.* **165**, 199 (1938).
 - [18] J.-P. Uzan, *Rev. Mod. Phys.* **75**, 403 (2003).
 - [19] G. A. Mena Marugán and S. Carneiro, *Phys. Rev. D* **65**, 087303 (2002).
 - [20] R. Bousso, *Rev. Mod. Phys.* **74**, 825 (2002); T. Jacobson and R. Parentani, *Found. Phys.* **33**, 323 (2003); T. Padmanabhan, gr-qc/0311036.
 - [21] By using (8), (11) and (18), it is possible to rewrite the solution (17) as $a = a_0[1 + nH_0(t - t_0)]^{1/n}$, where a_0 and H_0 are, respectively, the scale factor and the Hubble parameter at $t = t_0$. In the limit $n \rightarrow 0$, we obtain the de Sitter solution $a = a_0 \exp[H_0(t - t_0)]$, as should be.
 - [22] A. Dev, M. Sethi and D. Lohiya, *Phys. Lett. B* **504**, 207 (2001); D. Jair, A. Dev and J. S. Alcaniz, *Class. Quant. Grav.* **20**, 4163 (2003). See also M. V. John and J. V. Narlikar, *Phys. Rev. D* **65**, 043506 (2002).
 - [23] B. M. S. Hansen *et al.*, *Astrophys. J.* **574**, L155 (2002).
 - [24] M. Tegmark *et al.*, *Phys. Rev. D* **69**, 103501 (2004).
 - [25] In Planck units, equations (26)-(27) lead to $\Lambda \approx m^6$ (except for $n = 3$, in which case the cosmological constant is zero). This result is just the empirical relation originally considered by Zel'dovich, 37 years ago [7, 11].
 - [26] For $R = H^{-1} \approx \Lambda^{-1/2}$, we have $m_0 \approx \sqrt{\Lambda}$. Paul Weson (*Mod. Phys. Lett. A* **19**, 1995 (2004)) has recently proposed such a quantum of mass, on the basis of dimensional analysis and high dimensional theories.
 - [27] R. Penrose, *The Emperor's New Mind* (Oxford University Press, 1989), chapter 7.
 - [28] R. A. J. Matthews, *Astron. & Geophys.* **39**, 19 (1998).
 - [29] N. Banerjee and D. Pavón, *Class. Quant. Grav.* **18**, 593 (2001); *Phys. Rev. D* **63**, 043504 (2001).